A MOIE ON PROCESSING SEA LEVEL DATA(U) MAUAL POSTGRADUATE SCHOOL MONTEREY CA L C BREAKER 91 AUG 86 NPS-68-86-965 AD-A170 987 1/1 UNCLASSIFIED F/G 8/3 NL



MICROCOPY RESOLUTION TEST CHART NATIONAL BUREAU OF STANDARDS 1963 A



NPS68-86-005

NAVAL POSTGRADUATE SCHOOL

Monterey, Galifornia



A NOTE ON PROCESSING SEA LEVEL DATA

BY

LAURENCE C. BREAKER

JULY 1986

Approved for public release; distribution unlimited

Prepared for: Chief of Naval Research

Arlington, VA 22217



86 8 19 017

TIC FILE COPY

NAVAL POSTGRADUATE SCHOOL

Monterey, California 93943

RADM R.C. Austin Superintendent

David A. Schrady Provost

This report is for the research project "Ocean Prediction Through Observation, Modeling, and Analysis" sponsored by the Physical Oceanography Program of the Office of Naval Research under Program Element 61153N. Reproduction of all or part of this report is authorized.

This report was prepared by:

LAURENCE C. BREAKER
Adj. Prof. of Oceanography

CHRISTOPHER N.K. MOOERS
Professor and Chairman,
Department of Oceanography

Reviewed by:

Released by:

CHRISTOPHER N.K. MODERS, Chairman

Department of Oceanography

JOHN N. DYER

Dean of Science and Engineering

	REPORT DOCUMENTATION PAGE						
'S REPORT SECURITY CLASSIFICATION UNCLASSIFIED		D RESTRICTIVE MARKINGS					
2a SECURITY CLASSIFICATION AUTHORITY		3 DISTRIBUTION	3 DISTRIBUTION / AVAILABILITY OF REPORT				
2b DECLASSIFICATION / DOWNGRADING SCHEDU	LE	1	for public r	•	:		
			ion unlimite				
1 PERFORMING ORGANIZATION REPORT NUMBE	R(S)	5 MONITORING	ORGANIZATION R	EPORT NUMBER(5)		
NPS-68-86-005		1					
6a NAME OF PERFORMING ORGANIZATION	66 OFFICE SYMBOL	7a. NAME OF MONITORING ORGANIZATION					
NAVPGSCOL Dept. of Oceanography	(If applicable) 68						
6c. ADDRESS (City, State, and ZIP Code)	08	7b. ADDRESS (City, State, and ZIP Code)					
			,,	,			
Monterey, California 93943-50	00	1					
8a. NAME OF FUNDING / SPONSORING	86 OFFICE SYMBOL	9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER			UMBER		
ORGANIZATION	(If applicable)	1					
Office of Naval Research	(1122 PO)	N000146WR		né			
BC ADDRESS (City, State, and ZIP Code)		PROGRAM	PROJECT	TASK	WORK UNIT		
		ELEMENT NO	NO.	NO.	ACCESSION NO		
Arlington, VA 22217		61153N	RR0310306	<u> </u>			
The fire (include security classification)							
A Note on Processing Sea Level	Data						
12 PERSONAL AUTHOR(S)							
Laurence C. Breaker '3a TYPE OF REPORT 13b. TIME CO	DVERED	14. DATE OF REPO	OFT (Year Month	Dav) 15 PAGE	COUNT		
Technical FROM Ju	1 86 to Aug 86	1 August	1986	73 7202			
16 SUPPLEMENTARY NOTATION							
·							
17 COSATI CODES	18 SUBJECT TERMS (Continue on revers	e if necessary and	d identify by blo	ck number)		
F ELD GROUP SUB-GROUP							
	1						
'3 ABSTRACT (Continue on reverse if necessary	and identify by block	number)					
Procedures for processing hour	ly observations	of tidal ala	avation from	NOAA/NOS +	ido gougos		
for the study of subtidal vari	ability are out:	lined. Proce	essing proce	dures inclu	ide gauges ide an		
adjustment for datum reference	, low-pass filte	ering to remo	ove the diur	nal and sem	nidiurnal		
tides, and a static correction	for atmospheric	pressure.	These proce	dures are i	llustrated		
using a sequence of tidal elevents. September to 15 December 1984.	ation data trom	Santa Monica	a, Californi	a for the p	eriod 15		
September to 15 December 1964.							
20 DISTRIBUTION / AVAILABILITY OF ABSTRACT SOUNCLASSIFIED/UNLIMITED SAME AS RPT DTIC USERS		21 ABSTRACT SECURITY CLASSIFICATION					
MUNCLASSIFIED/UNLIMITED SAME AS RPT DTIC USERS 122 NAME OF RESPONSIBLE INDIVIDUAL		UNCLASSIFIED 22b TELEPHONE (Include Area Code) 22c OFFICE SYMBOL					

Laurence C. Breaker

A Note On Processing Sea Level Data

L.C. Breaker

Oceanography Department Naval Postgraduate School Monterey, CA 93943

Abstract. Procedures for processing hourly observations of tidal elevation from NOAA/NOS tide gauges for the study of subtidal variability are outlined. Processing procedures include an adjustment for datum reference, low-pass filtering to remove the diurnal and semidiurnal tides, and a static correction for atmospheric pressure. These procedures are illustrated using a sequence of tidal elevation data from Santa Monica, California for the period 15 September to 15 December 1984.

Accession For	
NTIS GRAGI	X
DTIC TAS Unamon mored	
Jutification.	
بالمسوعة أأأران والمساعلين والمساوية	
By	
Aunitalia	65
Dist	1
A-1	
<u> </u>	
	9110
	-ney)
Mar.	PECTED
1/2	

Introduction

The purpose of this note is to outline a set of procedures for processing hourly observations of tidal elevation to obtain information on subtidal variability.

Recommended Processing Procedures

Sea level data obtained from the National Ocean Survey of the National Oceanic and Atmospheric Administration (NOAA/NOS) tide gages is normally provided on an hourly basis with tidal elevations given in feet. The first correction to be applied adjusts the tidal elevations which are referenced to an arbitrary datum initially, to datums which are referenced to either Mean Higher High Water (MHHW), Mean High Water (MHW), Mean Sea Level (MSL), Mean Low Water (MLW), or Mean Lower Low Water (MLLW). NOAA/NOS provides the necessary data for each tide gauge location to make these corrections. The correction takes the following form:

Adjusted Sea Level (feet) = Raw Observation (feet) - Correction (feet)

To obtain MSL for example,

MSL (feet) = Raw Observations (feet) - MSL Correction (feet)

It is noted that because relative variations in sea level are generally of primary interest with regard to subtidal variability, no formal requirement to apply a datum correction in this case, exists.

The second adjustment that must be applied removes the diurnal and semidiurnal tidal components. The Godin low-pass digital filter is used to remove these components (Godin, 1972). The response of this filter can be expressed as

$$A_{25} \cdot A_{24}^{-2} / 25 \cdot 24^{2}$$

where

$$\tilde{A}_{n}(\sigma) = \frac{\sin(n\pi\Delta t\sigma)}{\sin(\pi\Delta\sigma\sigma)}$$

 σ = frequency in cycles per day

 Δt = time interval between observations in days


 A_n represents the arithmetric summation of n observations with the filter operating on the data three times, first using 25-point means (A_{25}) and then using 24-point means (A_{24}) , twice. According to Godin, the frequency band of this filter extends from 0 to 0.8 cycles per day (or 30 hours), beyond which the response is effectively zero. The response of this filter is shown in Figure 1. The half-amplitude cutoff is approximately 2.8 days (or ~66 hours). The amplitude of the diurnal and semidiurnal tidal components is reduced to less than 2% of the maximum value within the passband. A FORTRAN program implementing this tide-suppressing filter is contained in Appendix 1.

A third adjustment must be made to sea level data to remove the effects of atmospheric pressure. According to Lisitzin (1974), in the stationary case, the sea surface is depressed by 1 cm for every increase in atmospheric pressure of 1.005 mbar (and vice versa). It is noted, however, that such a simple static adjustment for atmospheric pressure does not take into account the effects of surface wind.

A relationship of the following form is recommended for making the sea level pressure (SLP) correction:

$$SL(t)_{adi} = (SL(t)-\overline{SL}) - (SLP(t)-\overline{SLP})$$
 (1)

* 1.000 vice 1.000cm/1.005 mbar is used here



igure 1. Response function for the Godin low-pass filter from hourly observations. The passband extends from 0 to ~66 hours.

where

 $SL(t)_{adj}$ = Pressure corrected sea level (cm) SL(t) = filtered sea level (cm) \overline{SL} = mean value of filtered sea level (cm) SLP(t) = filtered sea level pressure (mbar) \overline{SLP} = mean value of filtered sea level pressure (mbar)

The filtered sea level should first be converted to centimeters. Next, the SLP data should be low-pass filtered in a manner similar to that used for the sea level data. In this regard, it is recommended that the Godin filter be applied to the SLP data as well. Equivalent filtering of SLP has the advantage of producing a smooth time series with the same "time constant" or degree of smoothness as the sea level data. As indicated, both SL(t) and SLP(t) should be mean corrected in order to complete the correction for atmospheric pressure. Finally, whether or not the two terms on the R.H.S. of (1) are actually added or subtracted will depend on the sign convention used to define sea level and sea level pressure. The correct choice, of course, will produce an adjusted series with decreases (increases) in sea level that correspond to increases (decreases) in sea level pressure.

An Example

A 92-day sequence of hourly sea level data from the NOAA/NOS tide gauge at Santa Monica, California was acquired to illustrate the foregoing procedures. A datum correction furnished with the data from NOAA/NOS was applied initially to produce a sequence referenced to MSL. Next, the Godin low-pass filter was applied to the sea level data to remove the diurnal and


semidiurnal tidal components (Figure 2). The upper panel in Figure 2 shows the mixed character of this tidal record which clearly contains both diurnal and semidiurnal components. The lower panel shows the sea level record with the tides removed. To apply the correction for atmospheric pressure, hourly SLP data from a nearby NDBC environmental data buoy (NDBC Buoy 46011 off Pt. Conception) were acquired. These data were likewise smoothed using the Godin low-pass filter (Figure 3). The predominant diurnal component in SLP is effectively removed using this filter (lower trace). In the final figure (Figure 4), the mean-corrected SLP (upper panel) is subtracted from the mean-corrected sea level (middle panel) to yield sea level which has been corrected for atmospheric pressure (lower panel).

Discussion


Many different factors influence sea level. The time scales over which these factors are important vary from seconds to millenia. Only relatively short-term effects on sea level (< a few days) have been considered in the foregoing discussion. Although use of the Godin filter was recommended for removing the semidiurnal and diurnal tides, other tide-removing filters exist (Walters and Heston, 1982). In fact, Walters and Heston found that a so-called transform filter they describe generally had more desirable properties than the Godin filter. The only significant drawback they found in using the Godin filter is that in addition to removing tidal variations, it also attenuates variations in the 2 to 4 day range. For applications where information in this particular band is not of primary interest, however, the Godin filter is quite satisfactory.

The simple adjustment for atmospheric pressure described here assumes that the direct response of the ocean to a change in SLP is hydrostatic.


Hamon (1966), however, has shown that such a simple one-to-minus one correspondence between SLP and sea level may not always be observed. Factors such as coastal trapped waves may contribute to the deviations from the hydrostatic approximation (Hamon, 1976). It is also possible to examine the phase relationship between SLP and sea level using the techniques of cross-spectral analysis (Hamon, 1966). When such an analysis is conducted, corrections for atmospheric pressure can be defined for a range of frequencies. Using this approach, Hamon (1966) found that changes in sea level often follow changes in SLP with lags ranging from 0 to 2 days.



Hourly observations of sea level at Santa Monica, California from 15 September 1984. Upper panel shows the original, unfiltered data and the lower panel the low-passed data using the Godin filter. Figure 2.



Hourly observations of sea level pressure at NDBC Environmental Data Buoy 46011 (34.9N, 120.9W). The upper trace represents the original unfiltered data and the lower trace the low-passed data using the Godin filter. Figure 3.



Filtered sea level pressure (upper panel), filtered sea level (middle panel), and filtered sea level corrected for filtered sea level pressure (lower panel). Figure 4.

References

- Godin, G., 1972: The Analysis of Tides, University of Toronto Press, Toronto, 264 pp.
- Hamon, B.V., 1966: Continental shelf waves and the effects of atmospheric pressure and wind stress on sea level. Jour. Geophys. Res., 71, 2883-2893.
- Hamon, B.V., 1979: Generation of shelf waves on the East Australian coast by wind stress. Mem. Soc. Roy. des Sci. de Liege, 10, 359-367.
- Lisitzin, E., 1974: Sea Level Changes, Elsevier, Amsterdam, 286 pp.
- Walters, R.A. and C. Heston, 1982: Removing tidal-period variations from time-series data using low-pass digital filters. Jour. of Phys. Oceanogr., 12, 112-115.

```
FILE: GCDIN
                                                                                                                                                                                                                            FORIBAN
                                                                                          GODIN FILTER APPLIES A 24 HP BOTCAR LAICT AND A 25 HR BOTCAR CNCE TO HOURLY DATA. THE FIRST OUTPUT POINT IS 36 HOURS LATER THAN THE FIRST INPUT POINT.
 CCCC
                                CYCE TO HOUSELY DATA. THE FIRST COTEST IS 36 HOUSE LATER
INAN THE FIRST INSUT SCINT.

CIMENSION SL (3222)
INSUT ACTUAL VIMBER OF DATA ECINTS TO BE FRAD IN THE FOLLOWING
STATEMENT
NOTED = NUMBER OF INPUT DATA FOINTS

DO 66 J=1 NNEAD

REAC(5 11) SL (J)

COCRTINUS

11 FORMAT (211 FORMAT)

CALL BOXCAR (SL, NEBAD, 21, 1, NOUT1)

CALL BOXCAR (SL, NOUT1, 21, 1, NOUT1)

WHITE (01 1)

14 FORMAT (1X, NUMBER OF FILTEBED VALUES'//)

15 FORMAT (1X, NUMBER OF FILTEBED VALUES'//)

16 O7 Ret NOUT3

17 FORMAT (1X, F7.1)

END

STOR

STOR
 C
 C
C
```

Copy available to DTIC does not permit fully legible reproduction

INITIAL DISTRIBUTION LIST

1.	Naval Postgraduate School Department of Oceanography Monterey, CA 93943		
	Prof. Christopher N.K. Mooers Dr. Laurence C. Breaker LCDR J. Edward Johnson, USN Dr. Roland W. Garwood Dr. David C. Smith, IV Dr. Gordon Groves	33 1 1 1 1	
2.	Office of Naval Research (ONR) 800 N. Quincy St. Arlington, VA 22217		
	Dr. Thomas W. Spence Dr. Thomas B. Curtin Dr. Dennis Conlon	1 1 1	
3.	College of Oceanography Oregon State University Corvallis, OR 97331		
	Prof. Robert L. Smith Dr. Adriana Huyer	1 1	
4.	Commanding Officer Fleet Numerical Oceanography Center (FNOC) Monterey, CA 93943		
	Mr. R. Michael Clancy Mr. Ken Pollak Ms. Evelyn Hesse	1	l l
5.	Naval Ocean Research and Development Activity (NORDA) NSTL Station Bay St. Louis, MS 39525		
	Dr. Steve A. Piacsek Dr. Dana A. Thompson Dr. Harley C. Hurlburt Dr. Alexander Warn-Varnas	:	l l l
6.	Department of Oceanography University of Hawaii 2525 Correa Road Honolulu, HI 96822		
	Prof. Lorenz Magaard	•	1

7.	Applied Physics Laboratory University of Washington 1013 NE 40th Str. Seattle, WA 98105	
	Dr. Thomas B. Sanford	1
8.	School of Oceanography University of Washington Seattle, WA 98195	
	Dr. Steven C. Riser	1
9.	Pacific Marine Environmental Lab NOAA Bldg. 3 7600 Sand Point Way, NE Seattle, WA 98115	
	Mr. James R. Holbrook	1
10.	Naval Environmental Prediction Research Facility (NEPRF)	
	Ms. Marie Colton	1
11.	Allan Hancock Foundation University of Southern California Los Angeles, CA 90089-0371	
	Dr. Burton H. Jones	1
12.	Defense Technical Information Center Cameron Station Alexandria, VA 22314	2
13.	Oudley Knox Library Code 0142 Naval Postgraduate School Monterey, CA 93943	2
14.	Research Administration (Code 012) Naval Postgraduate School Monterey, CA 93943	1
15.	Moss Landing Marine Lab California State Colleges Sandholt Road Moss Landing, CA 95039	1
	Dr. John H. Martin	

16. NOAA/NMFS/Pacific Environmental Group P.O. Box 831 Monterey, CA 93942

Mr. Andy Bakun

